



Cannabis Information for Health Care Providers

Research shows that cannabis (marijuana, weed, pot, etc.) use during pregnancy or breastfeeding can have negative health impacts on developing fetuses and infants. There are also harmful impacts on the health and well-being of youth when cannabis is consumed during adolescence. Here are recommendations from professional organizations serving pregnant women, their infants, and children, as well as key data and information from scientific research on the health risks associated with cannabis use.



Using Cannabis During Pregnancy is Not Recommended

- The American College of Obstetricians and Gynecologists recommend women who are pregnant or contemplating pregnancy should be encouraged to discontinue use of cannabis.¹
- If a pregnant patient is already using cannabis for medicinal purposes, use should be discontinued in favor of an alternative treatment shown to be safe during pregnancy.¹
- Possible negative effects of using cannabis during pregnancy include fetal growth restriction and low birth weight.²
- Research has demonstrated that cannabis use may increase the chance of having a stillbirth.^{3,4}
- Evidence indicates cannabis use during pregnancy,

or while breastfeeding, can affect infant neurodevelopment.^{5,6}

Using Cannabis While Breastfeeding is Not Recommended

- The American Academy of Pediatrics and other professional organizations recommend that women do not use cannabis while breastfeeding.^{7,8}
- Regardless of the method of consumption (smoking, vaping, eating, or drinking), the active ingredient in cannabis, tetrahydrocannabinol (THC), is present in the breast milk of women who use cannabis and is transmitted to infants who nurse.^{9,10}
- Infants exposed to breast milk that contains THC may have trouble nursing because of sedation, reduced muscular tone, and poor sucking ability.¹¹

How Cannabis Affects Children and Youth

- The American Academy of Pediatrics recommends infants not be exposed to cannabis, given research demonstrating it may have negative effects on health and brain development.¹²
- Secondhand smoke from cannabis products entering the lungs of infants and children contains THC as well as many of the same chemicals as tobacco smoke.^{13,14}

- The way cannabis plants are grown has changed over the past few decades. Many plants now contain higher amounts of THC. The higher the THC content, the stronger the effects on your brain and behavior.^{15,16,17}
- Since the brain is still developing into the mid-20s, using cannabis during adolescence can negatively affect the developing brain, leading to short-term and long-term consequences. Such consequences may jeopardize educational, professional, and social achievements.¹⁸
- Research shows that youth who use cannabis before age 18, or use cannabis regularly, may be at higher risk of:
 - Skipping classes¹⁹
 - Lower grades and an increased risk of dropping out of school^{20,21}
 - Unemployment or having less fulfilling jobs later in life^{21,22}
- Cannabis use in adolescence can lead to addiction and increase the risk of use or abuse of other substances and illegal drugs.^{20,23}

Cannabis Use May Affect Youth Mental Health

- Regular cannabis use has been linked to anxiety, depression, suicidal thoughts and attempts, and death by suicide, especially for teens with a family history of mental illness.^{24,25,26}
- Although rare, cannabis use also increases the risk of schizophrenia and other psychoses, with the highest risk among those who use cannabis more frequently.^{27,28}

Communicate with Your Patients

- It is important to ask your patients about their cannabis use and provide information, without judgment, about safety concerns and referral for treatment if needed.

- More than 1 in 10 pregnant and non-pregnant women reported using cannabis in the past 12 months.²⁹
- In 2016, less than 1 in 5 high school students in California reported using cannabis in the past 30 days.³⁰
- Both groups reported beliefs that there is slight or no risk of harm from using cannabis once or twice a week.^{31,32}
- As acceptance of cannabis use increases, it is important to counsel pregnant women on the potential medical consequences of use during pregnancy.^{33,34}



LET'S TALK CANNABIS

 <http://bit.do/letstalkcannabis>
 letstalkcannabis@cdph.ca.gov

Under California law, adults 21 or older can use, carry, and grow cannabis (marijuana, weed, pot, etc.). Buying cannabis (without a valid physician's recommendation or a county-issued medical marijuana identification card) will become legal under California law for adults 21 or older on January 1, 2018. Use of medicinal cannabis is legal under California law if you have a valid physician's recommendation or a valid county-issued medical marijuana identification card. To buy medicinal cannabis, you must be 18 or older and have either a valid physician's recommendation, a valid county-issued medical marijuana identification card, or be a Primary Caregiver as defined in Health and Safety Code Section 11362.7(d) or 11362.5(e), with a valid physician's recommendation for the patient. In addition, consistent with the Compassionate Use Act, you may possess or cultivate any amount that is reasonably related to your current medical needs. The new California law, known as the Medicinal and Adult-Use Cannabis Regulation and Safety Act³⁵, includes information about where you can use cannabis, how much you can possess, and the penalties for illegal use. For more information, visit: https://leginfo.ca.gov/faces/billTextClient.xhtml?bill_id=20170180SB94.

References

- American College of Obstetricians and Gynecologists. 2015. "Committee Opinion No. 637: Marijuana Use During Pregnancy and Lactation." *Obstet Gynecol* 126 (1):234-8. doi: 10.1097/01.AOG.0000467192.89321.a6.
- Gunn, J. K., C. B. Rosales, K. E. Center, A. Nunez, S. J. Gibson, C. Christ, and J. E. Ehiri. 2016. "Prenatal exposure to cannabis and maternal and child health outcomes: a systematic review and meta-analysis." *BMJ Open* 6 (4):e009986. doi: 10.1136/bmjopen-2015-009986.
- Varner, M. W., R. M. Silver, C. J. Rowland Hogue, M. Willinger, C. B. Parker, V. R. Thorsten, R. L. Goldenberg, G. R. Saade, D. J. Dudley, D. Coustan, B. Stoll, R. Bukowski, M. A. Koch, D. Conway, H. Pinar, and U. M. Reddy. 2014. "Association between stillbirth and illicit drug use and smoking during pregnancy." *Obstet Gynecol* 123 (1):113-25. doi: 10.1097/aog.0000000000000052.
- National Institute of Health, Eunice Shriver Kennedy National Institute of Child and Human Development. 2013. "Tobacco, drug use in pregnancy can double risk of stillbirth." News Releases. December 11, 2013. <https://www.nih.gov/news-events/news-releases/tobacco-drug-use-pregnancy-can-double-risk-stillbirth>
- Wu, C. S., C. P. Jew, and H. C. Lu. 2011. "Lasting impacts of prenatal cannabis exposure and the role of endogenous cannabinoids in the developing brain." *Future Neurol* 6 (4):459-480.
- Leech, S. L., G. A. Richardson, L. Goldschmidt, and N. L. Day. 1999. "Prenatal substance exposure: effects on attention and impulsivity of 6-year-olds." *Neurotoxicol Teratol* 21 (2):109-18.
- American Academy of Pediatrics. 2012. "Section on Breastfeeding: Breastfeeding and the Use of Human Milk." *Pediatrics* 129 (3):e827-e841. doi: 10.1542/peds.2011-3552.
- Reece-Stremtan, S., and K. A. Marinelli. 2015. "ABM clinical protocol #21: guidelines for breastfeeding and substance use or substance use disorder, revised 2015." *Breastfeed Med* 10 (3):135-41. doi: 10.1089/bfm.2015.9992.
- Garry, A., V. Rigourd, A. Amirouche, V. Fauroux, S. Aubry, and R. Serreau. 2009. "Cannabis and breastfeeding." *J Toxicol* 2009:596149. doi: 10.1155/2009/596149.
- Perez-Reyes, M., and M. E. Wall. 1982. "Presence of delta9-tetrahydrocannabinol in human milk." *N Engl J Med* 307 (13):819-20. doi: 10.1056/nejm198209233071311.
- Liston, J. 1998. "Breastfeeding and the use of recreational drugs--alcohol, caffeine, nicotine and marijuana." *Breastfeed Rev* 6 (2):27-30.
- American Academy of Pediatrics. 2015. "The impact of marijuana policies on youth: clinical, research, and legal update." *Pediatrics* 135 (3):584-7. doi: 10.1542/peds.2014-4146.
- Herrmann, E. S., E. J. Cone, J. M. Mitchell, G. E. Bigelow, C. LoDico, R. Flegel, and R. Vandrey. 2015. "Non-smoker exposure to secondhand cannabis smoke II: Effect of room ventilation on the physiological, subjective, and behavioral/cognitive effects." *Drug Alcohol Depend* 151:194-202. doi: 10.1016/j.drugalcdep.2015.03.019.
- Moore, C., C. Coulter, D. Uges, J. Tuyay, S. van der Linde, A. van Leeuwen, M. Garnier, and J. Orbita, Jr. 2011. "Cannabinoids in oral fluid following passive exposure to marijuana smoke." *Forensic Sci Int* 212 (1-3):227-30. doi: 10.1016/j.forsciint.2011.06.019.
- ElSohly, M. A., Z. Mehmedic, S. Foster, C. Gon, S. Chandra, and J. C. Church. 2016. "Changes in Cannabis Potency Over the Last 2 Decades (1995-2014): Analysis of Current Data in the United States." *Biol Psychiatry* 79 (7):613-9. doi: 10.1016/j.biopsych.2016.01.004.
- Mehmedic, Z., S. Chandra, D. Slade, H. Denham, S. Foster, A. S. Patel, S. A. Ross, I. A. Khan, and M. A. ElSohly. 2010. "Potency trends of Delta9-THC and other cannabinoids in confiscated cannabis preparations from 1993 to 2008." *J Forensic Sci* 55 (5):1209-17. doi: 10.1111/j.1556-4029.2010.01441.x.
- National Institute on Drug Abuse. "Marijuana." Last updated April 2017. <https://d14rmgtrwzf5a.cloudfront.net/sites/default/files/1380-marijuana.pdf>
- National Academies of Sciences, Engineering, and Medicine. 2017. "The Health Effects of Cannabis and Cannabinoids: The Current State of Evidence and Recommendations for Research." Washington, DC: The National Academies Press. doi: 10.17226/24625.
- Arria, A. M., L. M. Garnier-Dykstra, E. T. Cook, K. M. Caldeira, K. B. Vincent, R. A. Baron, and K. E. O'Grady. 2013. "Drug use patterns in young adulthood and post-college employment." *Drug Alcohol Depend* 127 (1-3):23-30. doi: 10.1016/j.drugalcdep.2012.06.001.
- Macleod, J., R. Oakes, A. Copello, I. Crome, M. Egger, M. Hickman, T. Oppenkowski, H. Stokes-Lampard, and G. Davey Smith. 2004. "Psychological and social sequelae of cannabis and other illicit drug use by young people: a systematic review of longitudinal, general population studies." *Lancet* 363 (9421):1579-88. doi: 10.1016/s0140-6736(04)16200-4.
- Fergusson, D. M., and J. M. Boden. 2008. "Cannabis use and later life outcomes." *Addiction* 103 (6):969-76; discussion 977-8. doi: 10.1111/j.1360-0443.2008.02221.x.
- Zhang, C., J. S. Brook, C. G. Leukefeld, and D. W. Brook. 2016. "Trajectories of marijuana use from adolescence to adulthood as predictors of unemployment status in the early forties." *Am J Addict* 25 (3):203-9. doi: 10.1111/ajad.12361.
- Chen, C. Y., M. S. O'Brien, and J. C. Anthony. 2005. "Who becomes cannabis dependent soon after onset of use? Epidemiological evidence from the United States: 2000-2001." *Drug Alcohol Depend* 79 (1):11-22. doi: 10.1016/j.drugalcdep.2004.11.014.
- Kedzior, K. K., and L. T. Laefer. 2014. "A positive association between anxiety disorders and cannabis use or cannabis use disorders in the general population--a meta-analysis of 31 studies." *BMC Psychiatry* 14:136. doi: 10.1186/1471-244x-14-136.
- Lev-Ran, S., B. Le Foll, K. McKenzie, T. P. George, and J. Rehm. 2013. "Bipolar disorder and co-occurring cannabis use disorders: characteristics, co-morbidities and clinical correlates." *Psychiatry Res* 209 (3):459-65. doi: 10.1016/j.psychres.2012.12.014.
- Borges, G., C. L. Bagge, and R. Orozco. 2016. "A literature review and meta-analyses of cannabis use and suicidality." *J Affect Disord* 195:63-74. doi: 10.1016/j.jad.2016.02.007.
- Marconi, A., M. Di Forti, C. M. Lewis, R. M. Murray, and E. Vassos. 2016. "Meta-analysis of the Association Between the Level of Cannabis Use and Risk of Psychosis." *Schizophr Bull* 42 (5):1262-9. doi: 10.1093/schbul/sbw003.
- Di Forti, M., A. Marconi, E. Carra, S. Fraietta, A. Trotta, M. Bonomo, F. Bianconi, P. Gardner-Sood, J. O'Connor, M. Russo, S. A. Stilo, T. R. Marques, V. Mondelli, P. Dazzan, C. Pariante, A. S. David, F. Gaughran, Z. Atakan, C. Iyegbe, J. Powell, C. Morgan, M. Lynskey, and R. M. Murray. 2015. "Proportion of patients in south London with first-episode psychosis attributable to use of high potency cannabis: a case-control study." *Lancet Psychiatry* 2 (3):233-8. doi: 10.1016/s2215-0366(14)00117-5.
- Ko, J. Y., S. L. Farr, V. T. Tong, A. A. Creanga, and W. M. Callaghan. 2015. "Prevalence and patterns of marijuana use among pregnant and nonpregnant women of reproductive age." *Am J Obstet Gynecol* 213 (2):201.e1-201. e10. doi: 10.1016/j.ajog.2015.03.021.
- California Department of Public Health, Tobacco Control Program. 2016. "2016 California Student Tobacco Survey."
- Substance Abuse and Mental Health Services Administration. 2015. "Behavioral Health Barometer: California, 2015." U.S. Department of Health and Human Services. Rockville, MD: Substance Abuse and Mental Health Services Administration.
- Substance Abuse and Mental Health Services Administration. 2015. "2015 National Survey on Drug Use and Health." U.S. Department of Health and Human Services. Rockville, MD: Substance Abuse and Mental Health Services Administration.
- Harrison, Pam. "Marijuana Use Jumps in Pregnant Women." Medscape news release, December 29, 2016. <http://www.medscape.com/viewarticle/873834>.
- Phillips, Diana. "Few Ob/Gyns Counsel Pregnant Patients About Marijuana Use." Medscape news release, March 16, 2016. www.medscape.com/viewarticle/860456.
- Senate Bill No. 94, Sess. Of 2017 (Cal. 2017). https://leginfo.ca.gov/faces/billNavClient.xhtml?bill_id=201720180SB94